



MET3220C & MET6480 Computational Statistics



Parametric Probability Distributions

Discrete Distributions

Key Point: ALWAYS LOOK AT THE DATA!!!!
DOES THE DATA REALY FIT THE DISTRIBUTION

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability
Distributions 1

Parametric vs. Empirical Distributions

- **Empirical distributions** are based on a match to the sample data.
 - They are not based on underlying physical knowledge
- **Parametric distributions** are mathematical models (idealizations). In some cases the idealizations can be expected to be of very high quality.
 - One key question in later lectures will be 'how well does the distribution describe the data?'

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability
Distributions 2

Advantages of Parametric Distributions

- **Compactness:** The probability distribution of a parametric distribution can be explained with a formula.
 - Empirical distributions could require a very complicated histogram or pdf to describe the data.
- **Smoothing and interpolation:** Smoothing is largely unnecessary, or can be determined through integration of a formula. Interpolation is unnecessary.
 - For Empirical distributions, smoothing can be very complicated – or very misleading. Interpolation can be a nightmare in data sparse parts of the distribution.
- **Extrapolation:** If the parametric distribution is believed to be sufficiently accurate for conditions outside values in the sample data, then extrapolation is simply a matter of working with the equation describing the parametric distribution.

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability
Distributions 3

Discrete vs. Continuous Distributions

- Parametric distributions can be classified as either **discrete** or **continuous**.
- Continuous distributions apply to data that can have any value (including fractions) between specific limits.
 - Example: Gaussian distribution has limit of $\pm\infty$.
- Discrete distributions contain only specific values.
 - Example: a binomial distribution has integer values from 0 to N, where N is the number of samples.
 - E.g., five heads out of nine coin flips.

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability
Distributions 4

Binomial Distribution

- A binomial distribution describes the probability of occurrence of the number of times a outcome occurs, given a number of samples and the probability of that outcome for a single trial.
- For N trials, there will be N+1 possible outcomes, range from zero occurrences to N occurrences.
- Two key considerations for applicability of a binomial distribution are:
 - 1) the probability of occurrence does not change from event to event, and
 - 2) The outcome of each trial is independent from the other outcomes.
- Close approximations to these conditions are often acceptable.

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability
Distributions 5

Binomial Distribution Example

- Used to describe the outcome of a certain number of elementary events, for which the elementary event has only two possible outcomes.
 - Example: the daily maximum temperature $> 60^{\circ}\text{F}$
- These outcomes must be mutually exclusive.
- The events must be independent.
- The probability of the outcome of a single event must not change.
- Given N trials (N is the number of elementary events), there are N+1 possible outcomes.
 - E.g., 0, 1, 2, ..., N – 1, or N days when $T_{\max} > 60^{\circ}\text{F}$.
 - Note that for this example, the samples would have to be separated by quite a few days to truly be independent.

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability
Distributions 6

Binomial Distribution Formula

- Consider the probability of X of the N events occurring.
 - For example, X heads out of N coin flips.
- The probability of X events occurring ($\Pr\{X=x\}$) is written as

$$\Pr\{X=x\} = \binom{N}{x} p^x (1-p)^{N-x}, \text{ for } x = 1, 2, 3, \dots, N$$
- Where p is the probability of occurrence, and $(1-p)$ is the probability of non-occurrence.


$$\binom{N}{x} = \frac{N!}{x!(N-x)!}$$

Where $N! = 1 \times 2 \times 3 \times \dots \times N$

<http://campus.fsu.edu/~bourassa@met.fsu.edu>




Graphic from www.zoology.ubc.ca/~shl300/biomimetics.html



Parametric Probability Distributions 7

More Examples



Graphic from http://adept.maplesoft.com/categories/education/statistics/html/images/Binomial_Distribution/Binomial_Distribution29.gif


- Consider the following categories for each of the above plots:
 - $p < 0.5$, $p = 0.5$, $p > 0.5$
 - Identify the category for each plot.

<http://campus.fsu.edu/~bourassa@met.fsu.edu>



Parametric Probability Distributions 8

Example Binomial Distribution



- Note that the minimum in the peak occurs at $p = 0.5$.
- The spread of the distributions is largest for $p = 0.5$.

Graphic from <http://meted.ucar.edu/mwp/pcu1/ensemble/media/graphics/binomial.gif>



Parametric Probability Distributions 9

Binomial Distribution: Example

- The elementary event is the probability of a lake freezing in winter.
 - The probability in any year is 0.045
- Consider ten years of events.
- What is the probability of the lake freezing on at least one of those years?
 - $\Pr\{X \geq 1\} = \Pr\{X=1\} + \Pr\{X=2\} + \Pr\{X=3\} + \dots + \Pr\{X=10\}$
 - Or in an easier form: $\Pr\{X \geq 1\} = 1 - \Pr\{X=0\}$
 - $\Pr\{X \geq 1\} = 1 - \Pr\{X=0\} = 1 - [10!/(0!10!)](0.045)^0(0.955)^{10} = 0.37$

<http://campus.fsu.edu/~bourassa@met.fsu.edu>



Parametric Probability Distributions 10

Geometric Distribution

- The geometric distribution is used to determine the probability of a successful outcome in a fixed number of trials.
 - Otherwise the assumptions required to apply to the geometric distribution are identical to those required for a binomial distribution.
- The probability of a (one) successful trial in X events is
 - $\Pr\{X=x\} = p(1-p)^{x-1}$
- One application of this distribution is the length of waits or 'spells.' For example, wet spells or cold spells


<http://campus.fsu.edu/~bourassa@met.fsu.edu>



Parametric Probability Distributions 11


Example Geometric Distribution

$$p = 0.1$$



- Think about why there is a peak
 - Either probability of any success is small (left of peak), or
 - probability of a few successes is greater than probability of many successes.


<http://campus.fsu.edu/~bourassa@met.fsu.edu>



Parametric Probability Distributions 12

Negative Binomial Distribution

- The negative binomial distribution describes the probability of a number of failures (X), prior to the k^{th} success.
- In other words, after $X+k$ events, there were X failures in the first $X+k-1$ events. The last event must be the k^{th} success.
- $\Pr\{X=x\} = \Gamma(k+x) / (x! \Gamma(k)) p^k (1-p)^x$
- For typical applications k is an integer, in which case $\Gamma(k) = (k-1)!$



Graphic from <http://home.ubalt.edu/ntsbarsh/Business-stat/nebgi.jpg>


<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability Distributions 13

Example Negative Binomial Distribution



- The peak shifts to the right as the required number of successes increases. Why?

- Because more failures are likely to occur with the larger number of trials.

Graphic from <http://www.mathworks.com/access/helpdesk/help/toolbox/stats/nbncmpare.gif>

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability Distributions 14

Poisson Distribution

- The Poisson distribution describes the likelihood of a certain number of events (x) occurring in a limited space and/or time.
 - As with the previous distributions, the events must be independent.
- It is the limiting case of the binomial distribution, when $p \rightarrow 0$ and $N \rightarrow \infty$.
- Examples:
 - The number of hurricanes making landfall on the U.S. east coast in a year.
 - The annual number of tornadoes in Leon county
- The Poisson distribution is described by only one parameter: μ .
 - $\Pr\{X=x\} = \mu^x e^{-\mu} / x!$, for $x = 1, 2, 3, \dots$
 - This parameter is rather convenient because it is equal to both the standard deviation and the mean.


<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability Distributions 15


Example Poisson Distribution



- Poisson distributions are on the left, and the related cumulative probability distributions are on the right.
- Note that if you have a CDF, and a uniform random number generator, you can use the CDF to convert to the number associated with a randomly generated cumulative probability.

Graphic from http://en.wikipedia.org/wiki/Poisson_distribution

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability Distributions 16

Other Characteristics of the Discrete Parametric Parameterizations

- Expectation value
 - Expectation value is a fancy way of saying the mean or average.
 - Square brackets are usually used to indicate an expectation value.
 - For example $[x]$ (or $E[x]$) is the mean value of x .
- There are some simple rule that can be applied to products:
 - $E[c] = c$, where c is a constant
 - $E[c g(x)] = c E[g(x)]$, where $g(x)$ is a function of x .
 - $E[\sum g_i(x)] = \sum E[g_i(x)]$

Distribution	$\mu = E(X)$	$\sigma^2 = \text{Var}[X]$
Binomial	$N p$	$N p (1-p)$
Geometric	$1/p$	$(1-p) / p^2$
Negative Binomial	$k (1-p) / p$	$K (1-p) / p^2$
Poisson	μ	μ

<http://campus.fsu.edu/>
bourassa@met.fsu.edu



The Florida State University

Parametric Probability Distributions 17