



MET3220C & MET6480 Computational Statistics

Lecture 7

Parametric Probability Distributions

Discrete Distributions

**Key Point: ALWAYS LOOK AT THE DATA!!!
DOES THE DATA REALLY FIT THE DISTRIBUTION**



Programming Tidbit



- Several people have found that their correlate subroutine is returning a value of zero for correlation. They correctly point out that this does not make sense!
- The problem is related to the term $1 / (n - 1)$
- All terms in the equation are integers.
- For integer math, all fractions are truncated.
 - Therefore, $1 / (n - 1) = 0$ unless $n = 2$, which is not the case for assignment 3.



Parametric vs. Empirical Distributions



- **Empirical distributions** are based on a match to the sample data.
 - They are not based on underlying physical knowledge
- **Parametric distributions** are mathematical models (idealizations). In some cases the idealizations can be expected to be of very high quality.
 - One key question in later lectures will be ‘how well does the distribution describe the data?’



Advantages of Parametric Distributions



- **Compactness:** The probability distribution of a parametric distribution can be explained with a formula.
 - Empirical distributions could require a very complicated histogram or pdf to describe the data.
- **Smoothing and interpolation:** Smoothing is largely unnecessary, or can be determined through integration of a formula. Interpolation is unnecessary.
 - For Empirical distributions, smoothing can be very complicated – or very misleading. Interpolation can be a nightmare in data sparse parts of the distribution.
- **Extrapolation:** If the parametric distribution is believed to be sufficiently accurate for conditions outside values in the sample data, then extrapolation is simply a matter of working with the equation describing the parametric distribution.



Discrete vs. Continuous Distributions



- Parametric distributions can be classified as either **discrete** or **continuous**.
- Continuous distributions apply to data that can have any value (including fractions) between specific limits.
 - Example: Gaussian distribution has limit of $\pm\infty$.
- Discrete distributions contain only specific values.
 - Example: a binomial distribution has integer values from 0 to N, where N is the number of samples.
 - E.g., five heads out of nine coin flips.



Binomial Distribution



- A binomial distribution describes the probability of occurrence of the number of times a outcome occurs, given a number of samples and the probability of that outcome for a single trial.
- For N trials, there will be $N+1$ possible outcomes, range from zero occurrences to N occurrences.
- Two key considerations for applicability of a binomial distribution are:
 - 1) the probability of occurrence does not change from event to event, and
 - 2) The outcome of each trial is independent from the other outcomes.
- Close approximations to these conditions are often acceptable.



Binomial Distribution Example



- Used to describe the outcome of a certain number of elementary events, for which the elementary event has only two possible outcomes.
 - Example: the daily maximum temperature $> 60^{\circ}\text{F}$
- These outcomes must be mutually exclusive.
- The events must be independent.
- The probability of the outcome of a single event must not change.
- Given N trials (N is the number of elementary events), there are $N+1$ possible outcomes.
 - E.g., $0, 1, 2, \dots, N - 1$, or N days when $T_{\max} > 60^{\circ}\text{F}$.
 - Note that for this example, the samples would have to be separated by quite a few days to truly be independent.



Binomial Distribution Formula




- Consider the probability of X of the N events occurring.
- For example, X heads out of N coin flips.
- The probability of X events occurring ($\Pr\{X=x\}$) is written as

$$\Pr\{X = x\} = \binom{N}{x} p^x (1-p)^{N-x}, \text{ for } x = 1, 2, 3, \dots, N$$

- Where p is the probability of occurrence, and $(1-p)$ is the probability of non-occurrence.

$$\binom{N}{x} = \frac{N!}{x!(N-x)!}$$


Where $N! = 1 \times 2 \times 3 \times \dots \times N$




Graphic from www.zoology.ubc.ca/~bio300b/binomialnotes.html



More Examples



Graphic from www.phys.unsw.edu.au/~node13.html




Graphic from http://adept.maplesoft.com/categories/education/statistics/html/images/Binomial_Distribution/Binomial_Distribution29.gif

- Consider the following categories for each of the above plots:

- $p < 0.5$, $p = 0.5$, $p > 0.5$



Example Binomial Distribution



- Note that the minimum in the peak occurs at $p = 0.5$.
- The spread of the distributions is largest for $p = 0.5$.

Graphic from <http://meted.ucar.edu/nwp/pcu1/ensemble/media/graphics/binomial.gif>

<http://campus.fsu.edu/>
bourassa@met.fsu.edu

The Florida State University



Binomial Distribution: Example



- The elementary event is the probability of a lake freezing in winter.
- The probability in any year is 0.045
- Consider ten years of events.
- What is the probability of the lake freezing on at least one of those years?

$$\bullet \Pr\{X \geq 1\} = \Pr\{X=1\} + \Pr\{X=2\} + \Pr\{X=3\} + \dots + \Pr\{X=10\}$$

$$\bullet \text{ Or in an easier form: } \Pr\{X \geq 1\} = 1 - \Pr\{X=0\}$$

$$\bullet \Pr\{X \geq 1\} = 1 - \Pr\{X=0\} = 1 - [10! / (0!10!)] (0.045)^0 (0.955)^{10} \\ = 0.37$$



Geometric Distribution



- The geometric distribution is used to determine the probability of a successful outcome in a fixed number of trials.
 - Otherwise the assumptions required to apply to the geometric distribution are identical to those required for a binomial distribution.
- The probability of a (one) successful trial in X events is
- $\Pr\{X=x\} = p(1 - p)^{x-1}$
- One application of this distribution is the length of waits or ‘spells.’
For example, wet spells or cold spells




Example Geometric Distribution

$$p = 0.1$$



Graphics from http://zoonek2.free.fr/UNIX/48_R/g499.png



- Think about why there is a peak
- Either probability of any success is small (left), or
- probability of more than success is greater than probability of one success

<http://campus.fsu.edu/>

bourassa@met.fsu.edu

The Florida State University


Parametric Probability
Distributions 13



Negative Binomial Distribution



- The negative binomial distribution describes the probability of a number of failures (X), prior to the k^{th} success.
 - In other words, after $X+k$ events, there were X failures in the first $X+k-1$ events. The last event must be the k^{th} success.
- $\Pr\{X=x\} = \Gamma(k+x) / (x! \Gamma(k)) p^k (1 - p)^x$
- For typical applications k is an integer, in which case $\Gamma(k) = (k-1)!$




- Negative Binomial probability function with parameters $k = 6$, $x = 24$, $p = 0.5$

Graphic from <http://home.ubalt.edu/ntsbarsh/Business-stat/negbi.jpg>

<http://campus.fsu.edu/bourassa@met.fsu.edu>



Example Negative Binomial Distribution



- The peak shifts to the right as the required number of successes increases. Why?
- Because more failures are likely to occur with the larger number of trials.

<http://campus.fsu.edu/>

bourassa@met.fsu.edu

Graphic from <http://www.mathworks.com/access/helpdesk/help/toolbox/stats/nbincompare.gif>

Parametric Probability

The Florida State University

Distributions





Poisson Distribution



- The Poisson distribution describes the likelihood of a certain number of events (x) occurring in a limited space and/or time.
 - As with the previous distributions, the events must be independent.
- It is the limiting case of the binomial distribution, when $p \rightarrow 0$ and $N \rightarrow \infty$
- Examples:
 - The number of hurricanes making landfall on the U.S. east coast in a year.
 - The annual number of tornadoes in Leon county
- The Poisson distribution is described by only one parameter: μ .
 - $\Pr\{X=x\} = \mu^x e^{-\mu} / x!$, for $x = 1, 2, 3, \dots$
 - This parameter is rather convenient because it is equal to both the standard deviation and the mean.



Example Poisson Distribution



- Poisson distributions are on the left, and the related cumulative probability distributions are on the right.
- Note that if you have a CDF, and a uniform random number generator, you can use the CDF to convert to the number associated with a randomly generated cumulative probability.

Graphic from http://en.wikipedia.org/wiki/Poisson_distribution

<http://campus.fsu.edu/>
bourassa@met.fsu.edu

The Florida State University

Parametric Probability
Distributions 17



Other Characteristics of the Discrete Parametric Parameterizations



- Expectation value
 - Expectation value is a fancy way of saying the mean or average.
 - Square brackets are usually used to indicate an expectation value.
 - For example $E[x]$ (or $E[x]$) is the mean value of x .
- There are some simple rules that can be applied to products:
 - $E[c] = c$, where c is a constant
 - $E[c g(x)] = c E[g(x)]$, where $g(x)$ is a function of x .
 - $E[\sum g_j(x)] = \sum (E[g_j(x)])$

Distribution	$\mu = E(X)$	$\sigma^2 = \text{Var}[X]$
Binomial	Np	$N p (1 - p)$
Geometric	$1/p$	$(1 - p) / p^2$
Negative Binomial	$k (1 - p) / p$	$K (1 - p) / p^2$
Poisson	μ	μ